

March 1943



E-586

THE TOXICITY OF CERTAIN ORGANIC COMPOUNDS
TO YOUNG SCREWORM LARVAE

Roy Melvin, Raymond C. Bushland, and Charles L. Smith,
Division of Insects Affecting Man and Animals

The research program at the Menard, Tex., station includes the laboratory testing of materials for possible use as wound dressings to prevent infestation by larvae of the screwworm (Cochliomyia americana C. and P.). The method of testing and the results obtained with a large number of organic compounds have been reported by Bushland (1940a and b). The results obtained with 126 additional compounds supplied by the Division of Insecticide Investigations are presented in this paper.

As in the previous experiments, the desired amount of finely ground test material was mixed thoroughly with ground lean beef, bovine blood, water, and sufficient formaldehyde to retard putrefaction. Approximately 100 newly hatched larvae were placed on 150 grams of this treated medium in an uncovered wide-mouth pint Mason jar. The test jars were held at a temperature favorable for larval growth, and daily observations were made over a period of 3 days. The percentage mortality was then estimated, and any abnormalities in the surviving larvae were noted.

The larvae used in these tests were from the same inbred stock used in the work published by Bushland (1940a). Similar care was given in maintaining this stock to obtain uniform test larvae.

In these tests 0.25 gram of test material per jar (concentration 0.17 percent) was generally studied first. If all larvae were killed, in subsequent tests the dosage was reduced until some larvae survived. If all larvae were not killed at 0.25 gram per jar, the dosage was increased up to a maximum of 1.0 gram per jar (concentration 0.67 percent). All tests except those at the 1.0-gram dosage were conducted in duplicate, and with every lot of larvae four check jars were run. For these gross toxicity determinations it was not considered necessary to repeat tests, since experience had indicated that the results could generally be duplicated, and with toxic substances the results at dosages above and below the minimum lethal dosage served to confirm the observations.

The results of these tests are summarized in table 1. The various compounds are grouped according to their chemical classification. Two lethal concentrations are given, the highest concentration at which some larvae survived and the lowest concentration that caused 100 percent mortality (minimum lethal concentration) within the 3-day observation period. Materials described as slightly toxic are those failing to kill all larvae but causing some mortality or retardation in larval growth at a dosage of 1.0 gram per jar. Those compounds producing no noticeable ill effects at this dosage are designated as nontoxic.

Table 1.--Toxicity of various organic compounds against young screwworm larvae

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
Azo, hydrazo, and related compounds:		
p-Bromohydrazobenzene $\text{BrC}_6\text{H}_4\text{NHNHC}_6\text{H}_5$	0.05	0.08
p-Ethoxyazobenzene $\text{C}_6\text{H}_5\text{NNC}_6\text{H}_4\text{OC}_2\text{H}_5$.05	.08
p-Ethoxyhydrazobenzene $\text{C}_6\text{H}_5\text{NHNHC}_6\text{H}_4\text{OC}_2\text{H}_5$.08	.10
9-Fluorenonephenylhydrazone $\text{C}_6\text{H}_4\text{C}(\text{NNHC}_6\text{H}_5)\text{C}_6\text{H}_4$	Nontoxic	
p-Iodohydrazobenzene $\text{C}_6\text{H}_5\text{NHNHC}_6\text{H}_4\text{I}$.10	.17
p-Methylazobenzene $\text{C}_6\text{H}_5\text{NNC}_6\text{H}_4\text{CH}_3$.05	.08
p-Methylhydrazobenzene $\text{C}_6\text{H}_5\text{NHNHC}_6\text{H}_4\text{CH}_3$.08	.10
Heterocyclic compounds with sulfur outside ring:		
Piperidine-piperidyl dithio- carbamate $(\text{C}_5\text{H}_{10}\text{N})\text{C}(\text{S})\text{SH}(\text{C}_5\text{H}_{11}\text{N})$.17	.34
Thiocoumarin $\text{C}_6\text{H}_4\text{CH}=\text{CHC}(\text{S})\text{O}$	Less than .03	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
Heterocyclic compounds with 5-membered rings not containing sulfur:		
N-Benzylpyromucamide $(C_4H_3O)CONHCH_2C_6H_5$	0.05	0.08
N-Cyclohexylpyromucamide $(C_4H_3O)CONHC_6H_{11}$.05	.10
9-Furalfluorene $C_6H_4C [CH(C_4H_3O)] C_6H_4$.17	.67
2-Furanacrylamide $(C_4H_3O)CONH_2$.03	.05
beta-Furfuraldoxime $(C_4H_3O)CH(NO_2)$.10	.17
Phenazine oxide $C_6H_4NONC_6H_4$.01	.03
1,2,3,4-Tetrahydrocarbazole $C_6H_4NHC_6H_8$.17	.67
N-p-Tolylpyromucamide $(C_4H_3O)CONHC_6H_4CH_3$.08	.10
N-Xenylpyromucamide $(C_4H_3O)CONHC_6H_4C_6H_5$	Nontoxic	
N-(2,4-Xylyl)pyromucamide $(C_4H_3O)CONHC_6H_3(CH_3)_2$.10	.17

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	Percent	Percent
N-(2,5-Xylyl)pyromucamide (C ₄ H ₃ O)CONHC ₆ H ₃ (CH ₃) ₂	0.17	0.34
N-(2,6-Xylyl)pyromucamide (C ₄ H ₃ O)CONHC ₆ H ₃ (CH ₃) ₂	.03	.05
Heterocyclic compounds with 6-membered rings not containing sulfur:		
3-Acetyl-6-methyl-1,2-pyran-2, 4(3)dione (dehydroacetic acid) OCCHC(CH ₃)OC(O)CHCOCH ₃	.05	.10
2-Chloroacridone C ₆ H ₄ COC ₆ H ₃ ClNH	Nontoxic	
4-Chloroacridone C ₆ H ₄ COC ₆ H ₃ ClNH	Nontoxic	
2,3-Dihydroquinidine-12- carboxylic acid C ₆ H ₄ C(CO ₂ H)C(CH ₂) ₃ C=N	.17	.67
4-(2,4-Dinitrophenyl)morpholine (NO ₂) ₂ C ₆ H ₃ N(CH ₂ CH ₂) ₂ O	Nontoxic	
Dipiperonalcyclopentanone OCH ₂ OC ₆ H ₃ CHCC(O)C(CHC ₆ H ₃ OCH ₂ O)CH ₂ CH ₂	.17	.67
4-Hydroxyacridone C ₆ H ₄ COC ₆ H ₃ (OH)NH	.17	.67

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	Percent	Percent
Morpholine $\text{NH}(\text{CH}_2)_2\text{O}(\text{CH}_2)_2$	0.10	0.17
4-(4-Nitrophenyl)morpholine $\text{NO}_2\text{C}_6\text{H}_4\text{N}(\text{CH}_2\text{CH}_2)_2\text{O}$	Nontoxic	
2,2,6,6-Tetramethyl-1-nitroso-4-piperidone $(\text{CH}_3)_2\text{CCH}_2\text{COCH}_2\text{C}(\text{CH}_3)_2\text{NNO}$.17	.67
Xanthene $\text{C}_6\text{H}_4\text{CH}_2\text{C}_6\text{H}_4\text{O}$	Less than .03	
Xanthidrol $\text{C}_6\text{H}_4\text{CH}(\text{OH})\text{C}_6\text{H}_4\text{O}$.03	.05
Thiocyano compounds:		
p-Thiocyanobromobenzene $\text{BrC}_6\text{H}_4\text{SCN}$.05	.10
p-Thiocyaniodobenzene $\text{IC}_6\text{H}_4\text{SCN}$.10	.17
Amines, amides, imines, and imides containing sulfur:		
p-Acetamino-N-m-nitrophenylbenzene sulfonamide $\text{CH}_3\text{CONHC}_6\text{H}_4\text{SO}_2\text{NHC}_6\text{H}_4\text{NO}_2$	Nontoxic	
p-Acetamino-N-o-nitrophenylbenzene sulfonamide $\text{CH}_3\text{CONHC}_6\text{H}_4\text{SO}_2\text{NHC}_6\text{H}_4\text{NO}_2$	Nontoxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	Percent	Percent
p-Bromobenzenesulfonamide <chem>BrC6H4SO2NH2</chem>	0.01	0.03
p-Chlorobenzenesulfonamide <chem>ClC6H4SO2NH</chem>	.01	.03
Dithiooxamide <chem>NH2C(S)C(S)NH2</chem>	.08	.10
m-Nitrobenzenesulfonamide <chem>NO2C6H4SO2NH2</chem>	.03	.05
m-Nitro-N-phenylbenzenesulfonamide <chem>NO2C6H4SO2NHC6H5</chem>	Nontoxic	
3-Nitro-4-toluenesulfonamide <chem>CH3C6H3(NO2)SO2NH2</chem>	.10	.17
p-Toluenesulfonamide <chem>CH3C6H4SO2NH2</chem>	.10	.17
Sulfides, disulfides, and mercaptans:		
n-Butyl sulfone <chem>(C4H9)2SO2</chem>	.05	.08
Thio-2-naphthylmethyl ether <chem>C10H7SCH3</chem>	.10	.17
Aldehydes, ketones, thioaldehydes, and thioketones:		
Acenaphthenylphenyl ketone <chem>C6H5COC10H5(CH2)2</chem>	Nontoxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
Dianisalcyclopentanone $(\text{CH}_3\text{OC}_6\text{H}_4\text{CH})_2\text{CCH}_2\text{CH}_2\text{CCO}$	Slightly toxic	
Dibenzalcyclopentanone $\text{C}_6\text{H}_5\text{CH}=\text{CCH}_2\text{CH}_2\text{C}(=\text{CHC}_6\text{H}_5)\text{CO}$	Nontoxic	
alpha, alpha-Dichloro-p- phenylacetophenone $\text{C}_6\text{H}_5\text{C}_6\text{H}_4\text{COCHCl}_2$	0.10	0.17
2-Fluorylphenyl ketone $\text{C}_{13}\text{H}_9\text{COC}_6\text{H}_5$	Nontoxic	
p-Methylacetophenone $\text{CH}_3\text{C}_6\text{H}_4\text{COCH}_3$.10	.17
2-Nitrophenanthraquinone $\text{C}_6\text{H}_4\text{COCOC}_6\text{H}_3\text{NO}_2$	Nontoxic	
2,3-Octanedione-3-oxime $\text{CH}_3(\text{CH}_2)_4\text{C}(\text{NOH})\text{COCH}_3$	Slightly toxic	
Phenylglyoxal oxime $\text{C}_6\text{H}_5\text{COCHNOH}$.10	.17
1-Phenyl-1,2-propanedione-2- monoxime $\text{C}_6\text{H}_5\text{COC}(\text{NOH})\text{CH}_2\text{CH}_3$	Nontoxic	
2,6,7-Trinitrofluorenone $(\text{NO}_2)_2\text{C}_6\text{H}_2\text{COC}_6\text{H}_3\text{NO}_2$	Nontoxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
Phenols and phenol ethers:		
beta, beta'-Dinaphthol $\text{HO} \text{C}_{10}\text{H}_6 \text{C}_{10}\text{H}_6 \text{OH}$	Nontoxic	
3-Hydroxy-4-nitro-2-naphthoic acid $\text{C}_{10}\text{H}_5(\text{NO}_2)(\text{OH})\text{COOH}$	0.17	0.67
2-Nitroso-1-naphthol $\text{C}_{10}\text{H}_6(\text{NO})\text{OH}$	Slightly toxic	
Quinhydrone $\text{C}_6\text{H}_4\text{O}_2\text{C}_6\text{H}_4(\text{OH})_2$.17	.67
Amines, amides, imines, and imides not containing sulfur:		
Acetoacetic acid ethyl ester semicarbazone $\text{CH}_3\text{C}(\text{NNHCONH}_2)\text{CH}_2\text{COOC}_2\text{H}_5$	Slightly toxic	
Acetonyl acetone disemicarbazone $[\text{CH}_3\text{C}(\text{NNHCONH}_2)_2\text{CH}_2]_2$	Slightly toxic	
Acetophenone semicarbazone $\text{C}_6\text{H}_5\text{C}(\text{NNHCONH}_2)\text{CH}_3$.05	.10
alpha-Anisaldoxime $\text{CH}_3\text{OC}_6\text{H}_4\text{CH}=\text{NOH}$.10	.17
Benzaldehyde semicarbazone $\text{C}_6\text{H}_5\text{CHNNHCONH}_2$	Slightly toxic	
Benzoin semicarbazone $\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{C}(\text{NNHCONH}_2)\text{C}_6\text{H}_5$	Nontoxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
Benzophenone semicarbazone $(C_6H_5)_2CNNHCONH_2$	0.01	0.03
N-Carvacrylpalmitamide $C_{15}H_{31}CONHC_6H_3(CH_3)(C_3H_7)$	Nontoxic	
Carvacrylpropionamide $C_2H_5CONHC_6H_3(CH_3)(C_3H_7)$	Nontoxic	
omega-Chloroacetanilide $C_6H_5NHCCH_2Cl$	Less than .03	
p-Chloroacetophenone semicarbazone $ClC_6H_4C(CH_3)NNHCONH_2$	Nontoxic	
Cinnamaldehyde semicarbazone $C_6H_5CHCHCHNNHCONH_2$	Nontoxic	
Crotonaldehyde semicarbazone $CH_3CHCHCHNNHCONH_2$.05	.10
alpha-Cyanoacetanilide $CNCH_2CONHC_6H_5$	Slightly toxic 0.34	
Cyclopentanone semicarbazone $C_5H_8(NNHCONH_2)$.10	.17
Dibutylamine $(C_4H_9)_2NH$.03	.08
2,6-Dimethyl-4-heptanone semi- carbazone $[(CH_3)_2CHCH_2]_2CNNHCONH_2$.01	.03

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	Percent	Percent
Ethylmethyl ketone semicarbazone $C_2H_5C(NNHCONH_2)CH_3$	0.10	0.17
alpha-Ethylbutyraldehyde semi- carbazone $(C_2H_5)_2CHCHNNHCONH_2$.34	.67
9-Fluorenone oxime $C_6H_4C(=NOH)C_6H_4$	Nontoxic	
Fluorylamine $C_6H_4CH_2C_6H_3NH_2$.34	.67
2-Furaldehyde semicarbazone $(C_4H_3O)CHNNHCONH_2$.05	.10
p-Methylacetophenone semicarbazone $CH_3C_6H_4C(CH_3)NNHCOCH_3$	Nontoxic	
N-Methyl-m-acetotoluide $CH_3CON(CH_3)C_6H_4CH_3$.10	.17
4'-Methyl-4-nitrodiphenylamine $CH_3C_6H_4NHC_6H_4NO_2$	Nontoxic	
4-Methyl-2-pentanone semicarbazone $(CH_3)_2CHCH_2C(CH_3)NNHCONH_2$.17	.34
Methylpropyl ketone semicarbazone $CH_3C(NNHCONH_2)C_3H_7$.10	.17
p-Nitrosodiethylaniline $(C_2H_5)_2NC_6H_4NO$.08	.10
2-Octanone semicarbazone $CH_3(CH_2)_5C(CH_3)NNHCONH_2$	Slightly toxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
Phthalonitrile $C_6H_4(CN)_2$	0.03	0.05
Salicylaldehyde semicarbazone $HOC_6H_4CHNNHCONH_2$	Nontoxic	
2,4,2',4'-Tetrabromodiphenylamine $Br_2C_6H_3NHC_6H_3Br_2$	Nontoxic	
2,4,2',4'-Tetrachlorodiphenylamine $Cl_2C_6H_3NHC_6H_3Cl_2$	Nontoxic	
p-Tolyl lactamide $CH_3CHOHCONHC_6H_4CH_3$.17	.67
Bis-(2,4,6-trinitrophenyl)amine $[(NO_2)_3C_6H_2]_2NH$	Less than .03	
N-Xenylacetamide $CH_3CONHC_6H_4C_6H_5$	Slightly toxic	
N-Xenylbenzamide $C_6H_5CONHC_6H_4C_6H_5$	Nontoxic	
N-Xenylformamide $HCONHC_6H_4C_6H_5$	Nontoxic	
N-Xenylpropionamide $C_2H_5CONHC_6H_4C_6H_5$	Nontoxic	
N-(2,4-Xylyl)benzamide $C_6H_5CONHC_6H_3(CH_3)_2$	Nontoxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
N-(2,5-Xylyl)benzamide $C_6H_5CONHC_6H_3(CH_3)_2$	Nontoxic	
N-(2,6-Xylyl)benzamide $C_6H_5CONHC_6H_3(CH_3)_2$	Nontoxic	
N-(2,4-Xylyl)formamide $HCONHC_6H_3(CH_3)_2$	0.10	0.17
N-(2,5-Xylyl)formamide $HCONHC_6H_3(CH_3)_2$.05	.10
N-(2,6-Xylyl)formamide $HCONHC_6H_3(CH_3)_2$.05	.10
N-(2,4-Xylyl)propionamide $C_2H_5CONHC_6H_3(CH_3)_2$	Slightly toxic	
N-(2,5-Xylyl)propionamide $C_2H_5CONHC_6H_3(CH_3)_2$	Nontoxic	
N-(2,6-Xylyl)propionamide $C_2H_5CONHC_6H_3(CH_3)_2$.17	.34
Acids, alcohols, and esters:		
2-Acetoxy-2-naphthoic acid $CH_3COOC_{10}H_6COOH$.17	.67
2,4-Dinitrophenol acetate $CH_3CO_2C_6H_3(NO_2)_2$.05	.10
9-Fluorenol acetate $C_6H_4CH(OCOCH_3)C_6H_4$	Nontoxic	

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :
	<u>Percent</u> <u>Percent</u>
p-Nitroiodosobenzene acetate (NO ₂)C ₆ H ₄ I=(OCOCH ₃) ₂	Nontoxic
Nitro and nitroso compounds:	
alpha, beta-Dibromo-beta-nitroethylbenzene C ₆ H ₅ CHBrCHBrNO ₂	0.05 0.08
o-Iodosonitrobenzene C ₆ H ₄ (NO ₂)IO	.05 .08
1-Nitronaphthalene C ₁₀ H ₇ NO ₂	.05 .08
o-Nitronitrosobenzene NO ₂ C ₆ H ₄ NO	Nontoxic
beta-Nitrosostyrene C ₆ H ₅ CH=CHNO	.10 .17
4-Nitrosothymol (CH ₃) ₂ CHC ₆ H ₂ (CH ₃)(OH)NO	Nontoxic
2,4,6-Trinitrotoluene C ₆ H ₂ (CH ₃)(NO ₂) ₃	.08 .10
Aryl, alkyl, and aralkyl halides:	
3-Bromoacenaphthene C ₁₀ H ₅ (CH ₂) ₂ Br	.17 .67
2-Bromofluorene C ₆ H ₄ CH ₂ C ₆ H ₃ Br	.05 .08

Table 1 (Continued)

Compound	: Highest : : concen- : Minimum : tration : lethal : at which : concen- : larvae : tration : survived :	
	<u>Percent</u>	<u>Percent</u>
9-Chlorofluorene $\text{C}_6\text{H}_4\text{CHClC}_6\text{H}_4$	Nontoxic	
1-Chloronaphthalene $\text{C}_{10}\text{H}_7\text{Cl}$	0.10	0.17
Iodosobenzene $\text{C}_6\text{H}_5\text{IO}$.10	.17
2-Iodofluorene $\text{C}_6\text{H}_4\text{CH}_2\text{C}_6\text{H}_3\text{I}$	Nontoxic	

Materials that were found to be at least as toxic as diphenylamine 1/ are tabulated below. Compounds of similar effectiveness

1/ Bushland (1940a) reported that the minimum lethal concentration for diphenylamine was 0.10-0.17 percent (all larvae killed at 0.17 but some survived at 0.10 percent). Melvin and coworkers (1939a and b, 1941) have shown that diphenylamine is a practical screwworm remedy. Although in laboratory tests many compounds have been effective at lower dosages than are required for diphenylamine, field tests evaluating such qualities as duration of protection, toxicity to infested animals, and rate of healing of treated wounds have shown that the diphenylamine treatment is preferable to any others so far considered.

are listed alphabetically within groups, and the groups are arranged in decreasing order of toxicity.

Killed all larvae at a concentration of 0.03 percent:

Benzophenone semicarbazone
Bis-(2,4,6-trinitrophenyl)amine
p-Bromobenzenesulfonamide
omega-Chloroacetanilide
p-Chlorobenzenesulfonamide
2,6-Dimethyl-4-heptanone semicarbazone
Phenazine oxide
Thiocoumarin
Xanthene

Killed all larvae at a concentration of 0.05 percent but some survived
at 0.03 percent:

2-Furanacrylamide
m-Nitrobenzenesulfonamide
Phthalonitrile
Xanthydrol
N-(2,6-Xylyl)pyromucamide

Killed all larvae at 0.10 percent but some survived at 0.05 percent:

Acetophenone semicarbazone

3-Acetyl-6-methyl-1,2-pyran-2,4(3)-dione

(dehydroacetic acid)

N-Benzylpyromucamide

2-Bromofluorene

p-Bromohydrazobenzene

n-Butyl sulfone

Crotonaldehyde semicarbazone

alpha, beta-Dibromo-beta-nitroethylbenzene

Dibutylamine

2,4-Dinitrophenol acetate

Dithiooxamide

p-Ethoxyazobenzene

p-Ethoxyhydrazobenzene

2-Furaldehyde semicarbazone

o-Iodosonitrobenzene

p-Methylazobenzene

p-Methylhydrazobenzene

1-Nitronaphthalene

p-Nitrosodiethylaniline

p-Thiocyanobromobenzene

N-p-Tolylpyromucamide

2,4,6-Trinitrotoluene

N-(2,5-Xylyl)formamide

N-(2,6-Xylyl)formamide

Killed all larvae at 0.10 percent but some survived at 0.17 percent:

alpha-Anisaldoxime

1-Chloronaphthalene

Cyclopentanone semicarbazone

alpha, alpha-Dichloro-p-phenylacetophenone

Ethylmethyl ketone semicarbazone

beta-Furfuraldoxime

p-Iodohydrazobenzene

Iodosobenzene

p-Methylacetophenone

N-Methyl-m-acetotoluide

Methylpropyl ketone semicarbazone

Morpholine

beta-Nitrostyrene

3-Nitro-4-toluenesulfonamide

Phenylglyoxal oxime

p-Thiocyaniodobenzene

Thio-2-naphthylmethyl ether

p-Toluenesulfonamide

N-(2,4-Xylyl)formamide

N-(2,4-Xylyl)pyrrolisamide

SUMMARY

As a means of selecting larvicidal materials effective against the screwworm, 126 organic compounds were tested in the laboratory by the jar method. In these tests 37 compounds were more toxic than diphenylamine and 20 compounds were of about the same toxicity. Toxic materials were found in all 13 classes of organic compounds tested, and 40 nontoxic materials were present in 10 classes.

Newly hatched screwworm larvae failed to survive for 3 days when their food contained as much as 0.03 percent of any of the following compounds: bis-(2,4,6-Trinitrophenylamine), p-bromobenzenesulfonamide, omega-chloroacetanilide, p-chlorobenzenesulfonamide, 2,6-dimethyl-4-heptanone semicarbazone, phenazine oxide, thiocoumarin, and xanthene.

LITERATURE CITED

Bushland, R. C.

- 1940a. The toxicity of phenothiazine and certain related compounds to young screwworms. Jour. Econ. Ent. 33:666-9.
-

- 1940b. The toxicity of some organic compounds to young screwworms. Jour. Econ. Ent. 33:669-76.

Melvin, Roy, H. E. Parish, E. F. Knipling, and R. C. Bushland.

- 1939a. Results of studies on diphenylamine as a wound protector against the screwworm, Cochliomyia americana C. and P. U. S. Bur. Ent. and Plant Quar. E-480, 4 pp. [Processed.]
-

- 1939b. Diphenylamine as a wound protector against the screwworm, Cochliomyia americana C. and P. U. S. Bur. Ent. and Plant Quar. E-481, 4 pp. [Processed.]

Melvin, Roy, C. L. Smith, H. E. Parish, and W. L. Barrett, Jr.

1941. A new remedy for the prevention and treatment of screwworm infestations of livestock. U. S. Bur. Ent. and Plant Quar. E-540, 4 pp. [Processed.]

UNIVERSITY OF FLORIDA



3 1262 09224 7062